

Complementarity and the Afshar Experiment

Tabish Qureshi*

Department of Physics, Jamia Millia Islamia, New Delhi-110025, India.

A modified two-slit interference experiment, proposed and carried out by Shahriar Afshar [first reported by M. Chown, *New Scientist* **183**, 30 (2004)], claims to demonstrate a violation of Bohr's complementarity principle. We point out the flaw in Afshar's analysis and show why complementarity is robust and cannot be violated in any experiment which is a variant of the conventional two-slit experiment.

PACS numbers: 03.65.Ud ; 03.65.Ta

I. INTRODUCTION

It is a well established experimental fact that the wave and particle aspects of quantum systems are mutually exclusive. Niels Bohr elevated this concept, which is probably born out of the uncertainty principle, to the status of a separate principle, the principle of complementarity¹.

Since then, the complementarity principle has been demonstrated in various experiments, the most common of which is the two-slit interference experiment. It has been shown that if the which-way information in a double-slit experiment is stored somewhere, the interference pattern is destroyed, and if one chooses to "erase" the which-way information *after* detecting the particle, the interference pattern comes back. This phenomenon goes by the name of quantum erasure^{2,3}.

Shahriar Afshar proposed and carried out an interesting experiment which claims to violate the complementarity principle^{4,5}. A schematic diagram of the experiment is shown in FIG. 1. Basically, it consists of a standard two-slit experiment, with a converging lens L behind the conventional screen for obtaining the interference pattern. Although Afshar uses pinholes instead of slits, we will continue to refer to them as slits. If the screen is removed, the light passes through the lens and produces two images of the slits, which are captured on two detectors D_A and D_B respectively. Opening only

slit A results in only detector D_A clicking, and opening only slit B leads to only D_B clicking. Afshar argues that the detectors D_A and D_B yield information about which slit, A or B , the particle initially passed through. If one places a screen before the lens, the interference pattern is visible.

Conventionally, if one tries to observe the interference pattern, one cannot get the which-way information. Afshar has a clever scheme for establishing the existence of the interference pattern without actually observing it. First the exact location of the dark fringes are noted by observing the interference pattern. Then, thin wires are placed in the exact locations of the dark fringes. The argument is that if the interference pattern exists, sliding in wires through the dark fringes will not affect the intensity of light on the two detectors. If the interference pattern is not there, some photons are bound to hit the wires, and get scattered, thus reducing the photon count at the two detectors. This way, the existence of the interference pattern can be established without actually disturbing the photons in any way. This is similar in spirit to the so called "interaction-free measurements" where the non-observation of a particle along one path establishes that it followed the other possible path, without actually measuring it⁶. Afshar carried out the experiment and found that sliding in wires in the expected locations of the dark fringes, doesn't lead to any significant reduction of intensity at the detectors. Hence Afshar claims that he has demonstrated a violation of complementarity.

The complementarity principle is at the heart of quantum mechanics, and its violation will be deeply disturbing to its established understanding. As expected, there has been skepticism towards Afshar's experiment, and a heated debate is currently going on^{7,8,9,10,11}. However, none of the criticisms has been satisfactorily able to point out any flaw in Afshar's interpretation of the experiment. Most agree that if the introduction of wires has no effect on the intensity, it shows that interference exists. And everybody seems to agree that detecting a photon at (say) D_A means that it came from slit A . Most have fallen back to a more formal interpretation of Bohr's principle, namely that the wave and particle nature cannot be seen in the same experiment, for the same photons. They argue that Afshar has to do not one, but two experiments to prove his point - one without the wires, one with the




FIG. 1: A schematic representation of Afshar's experiment. Light emerges from two pin-holes (not slits) A and B and interferes. Thin wires are placed carefully in the exact locations of the dark fringes of the interference pattern. The lens L collects the light and obtains the images of the two slits A' and B' , respectively. The detectors D_A and D_B collect the photons for these two images.

wires. Some argue that Afshar already assumes the existence of fringes, and that his argument is circular. However, a reader who respects empirical facts, doesn't see it as two experiments if putting in the wires is not changing the results. One would say, the interference is out there in the middle, and one check it out that the photons are not passing through certain regions, the dark fringes.

Let us understand what is happening in the experiment slightly better by simplifying the experiment. One might argue that a two-slit experiment is the simplest experiment one can imagine. But this experiment still has a large number of degrees of freedom, and the Hilbert space is big. We will show in the following that the simplest interference (thought) experiment can be just carried out using a spin-1/2 particle.

II. A SIMPLIFIED "TWO-SLIT" EXPERIMENT

Let there be a spin-1/2 particle traveling along x-axis. For our purpose, its physical motion is unimportant - we will only be interested in the dynamics of its spin. In the two-slit experiment, the particle emerges from the slits in a superposition of two physically separated, localized wave-packets, which are orthogonal to each other. Their physical separation guarantees their orthogonality. Any subsequent unitary evolution will retain their orthogonality. We assume the initial state of the spin to be

$$|\psi\rangle_0 = \frac{1}{\sqrt{2}}(|\uparrow\rangle + |\downarrow\rangle). \quad (1)$$

Here, the states $|\uparrow\rangle$ and $|\downarrow\rangle$ are the eigenstates of $\hat{S}_z$, and play the role of the two wave-packets that emerge from the double-slit. The time evolution, in the conventional two-slit experiment, spreads the wave-packets so that they overlap. In our thought experiment, we employ a homogeneous magnetic field B , acting along the y-axis, to evolve the two states $|\uparrow\rangle$ and $|\downarrow\rangle$. Thus the Hamiltonian of the system is $\hat{H} = B\hat{S}_y$.

The time evolution operator can be written as

$$\begin{aligned} \hat{U}(t) &= \exp\left(\frac{i}{\hbar} B \hat{S}_y t\right) \\ &= \cos(Bt/2) + i\hat{\sigma}_y \sin(Bt/2), \end{aligned} \quad (2)$$

where $\hat{\sigma}_y$ is the usual Pauli matrix. The time evolution operator, for a time $\tau = \pi/2B$ has the form

$$\hat{U}(\tau) = \frac{1}{\sqrt{2}}(1 + i\hat{\sigma}_y). \quad (3)$$

It is straight forward to see that the time evolution transforms the states $|\uparrow\rangle$ and $|\downarrow\rangle$ as

$$\begin{aligned} \hat{U}(\tau)|\uparrow\rangle &= \frac{1}{\sqrt{2}}(|\uparrow\rangle - |\downarrow\rangle) \\ \hat{U}(\tau)|\downarrow\rangle &= \frac{1}{\sqrt{2}}(|\uparrow\rangle + |\downarrow\rangle) \end{aligned} \quad (4)$$

A further evolution through a time τ will transform the states as

$$\begin{aligned} \hat{U}(2\tau)|\uparrow\rangle &= -|\downarrow\rangle \\ \hat{U}(2\tau)|\downarrow\rangle &= |\uparrow\rangle \end{aligned} \quad (5)$$

After an evolution through a total time 2τ , if one puts a spin detector, one will either get a "spin-down" or a "spin-up". In the beginning, if we started out with a state $|\uparrow\rangle$, the detector at the end will register a $|\downarrow\rangle$ state. On the other hand, if we started with a $|\downarrow\rangle$ state, the detector at the end will register a $|\uparrow\rangle$ state. Thus, the detector at the end obtains a which-initial-state information about the spin, exactly as the detectors in Afshar experiment obtain a which-slit information.

So, now we carry out our thought experiment, with the initial state $|\psi\rangle_0$, as given by (10). After a time τ we get into a state which is equivalent to the interference region in Afshar's experiment:

$$\hat{U}(\tau)|\psi\rangle_0 = \frac{1}{2}(|\uparrow\rangle - |\downarrow\rangle) + \frac{1}{2}(|\uparrow\rangle + |\downarrow\rangle). \quad (6)$$

Equation (6) represents an interference pattern, because the "down-spins" cancel out to give destructive interference, while the "up-spins" add up to give constructive interference. So, in our two-state interference experiment, there is one dark fringe and one bright fringe. After parts of the states have been destroyed by the "dark-fringe", what is left is just

$$\hat{U}(\tau)|\psi\rangle_0 = \frac{1}{2}|\uparrow\rangle + \frac{1}{2}|\uparrow\rangle. \quad (7)$$

Now, these two contribution from the two initially orthogonal states are identical. States which are not orthogonal, are naturally not distinguishable. Thus the which-way or which-initial-state information is lost at this stage. It might serve some useful purpose by keeping the two contributions separate and evolving them for a further time τ to see what they yield at the spin detector.

$$\hat{U}(\tau)\left(\frac{1}{2}|\uparrow\rangle + \frac{1}{2}|\uparrow\rangle\right) = \frac{1}{2\sqrt{2}}(|\uparrow\rangle - |\downarrow\rangle) + \frac{1}{2\sqrt{2}}(|\uparrow\rangle - |\downarrow\rangle). \quad (8)$$

So, we see that both the initially orthogonal parts of the initial state lead to a superposition of $|\downarrow\rangle$ and $|\uparrow\rangle$ states. So, although the spin-detector at the end still gives either a "spin-up" or a "spin-down", it gives no information about whether the initial state was a $|\downarrow\rangle$ or $|\uparrow\rangle$.

III. INTERFERENCE DESTROYS WHICH-WAY INFORMATION

It is obvious from the simple thought experiment discussed above that interference destroys the which-way information. One might wonder if this statement is general enough, or if it is an artifact of the over-simplification of the analysis. Let us discuss this point in a bit more

detail. In any variant of the conventional two-slit experiment, the initial state has to be in a superposition of two orthogonal states. This follows simply from the fact that the two slits are distinguishable. Thus the initial state $|\psi\rangle_0$ could be written as $|\psi\rangle_0 = (|\psi_A\rangle + |\psi_B\rangle)/\sqrt{2}$. These states then evolve and reach the region where they interfere. The time evolution, being unitary, retains their orthogonality. Let us assume that in the region where they interfere, they have the form:

$$\begin{aligned} |\psi_A\rangle &= a|\alpha\rangle + c_1|\gamma\rangle \\ |\psi_B\rangle &= b|\beta\rangle - c_2|\gamma\rangle, \end{aligned} \quad (9)$$

where we assume that c_1 and c_2 are real and positive, although this requirement is not essential. Here, $|\alpha\rangle$, $|\beta\rangle$ and $|\gamma\rangle$ need not be simple states - each may involve a multitude of degrees of freedom. However, $|\psi_A\rangle$ and $|\psi_B\rangle$ have to have this form, so that there are parts from the two which cancel out, to give the so-called dark-fringes. We assume that $|\gamma\rangle$ is orthogonal to both $|\alpha\rangle$ and $|\beta\rangle$, for the dark fringe to be distinguishable from the rest of the part.

If complementarity could indeed be violated, the parts of the two initial states which are left (after the destructive interference), should be orthogonal, so that they can contain a which-way information. In our example, it would mean that the states $|\alpha\rangle$ and $|\beta\rangle$ should be orthogonal. It is trivial to verify that if $|\psi_A\rangle$ and $|\psi_B\rangle$ are orthogonal, $|\alpha\rangle$ and $|\beta\rangle$ can never be orthogonal. This general analysis shows that if interference exists, it necessarily destroys which-way information. So, complementarity is robust, and cannot be violated in any such interference experiment.

IV. REAL TWO-SLIT EXPERIMENT

Coming back to Afshar's experiment, let us see what implication the preceding analysis has on it. Clearly, in Afshar's experiment, after the particle pass through the interference region, the which way information is lost. The detector D_A clicking doesn't mean that the particle came from slit A . It might appear hard to visualize that a wave-packet which travels in a straight line from slit A can yield a click on detector D_B , which is not in its direct path. However, the argument of momentum conservation will hold only if we *knew* that the particle started out from slit A - in that situation it would never reach detector D_B . In order to demonstrate the validity of these arguments for Afshar's experiment, let us look at the two-slit experiment in more detail.

We carry out the analysis for massive particles, to show that the argument doesn't just hold for electromagnetic waves. Consider the particle to be moving along the x-axis, and the slit plane to be parallel to the y-axis. We will only be interested in the dynamics of the state in the y-direction, whereas the x-axis motion just serves the purpose of transporting the particle from the slits

to the detectors. Let us assume that when the particle emerges from the double-slit, its state is given by a superposition of two distinct, spatially localized wave-packets. The state at this time, which we choose to call $t = 0$, can be written as

$$\psi(y, 0) = \frac{a}{(\pi/2)^{1/4}\sqrt{\epsilon}} e^{-(y-y_0)^2/\epsilon^2} + \frac{b}{(\pi/2)^{1/4}\sqrt{\epsilon}} e^{-(y+y_0)^2/\epsilon^2} \quad (10)$$

where ϵ is the width of the wave-packets, $2y_0$ is the slit separation, and a and b are the amplitudes of the two wave-packets.

The wave-packets evolve in time, during which they spread, and reach the region where they overlap, and thus, interfere. This state can just be obtained by evolving (10), using the Hamiltonian $\hat{H} = \hat{p}_y^2/2m$. Hence, the state at time t can be written as

$$\psi(y, t) = aC(t)e^{-\frac{(y-y_0)^2}{\epsilon^2 + 2i\hbar t/m}} + bC(t)e^{-\frac{(y+y_0)^2}{\epsilon^2 + 2i\hbar t/m}} \quad (11)$$

where

$$C(t) = \frac{1}{(\pi/2)^{1/4}\sqrt{\epsilon + 2i\hbar t/m\epsilon}}. \quad (12)$$

Now, this state can also be rewritten as

$$\begin{aligned} \psi(y, t) &= aC(t)e^{-\frac{y^2+y_0^2}{\Omega(t)}} \left(\cosh\left(\frac{2yy_0}{\Omega(t)}\right) + \sinh\left(\frac{2yy_0}{\Omega(t)}\right) \right) \\ &+ bC(t)e^{-\frac{y^2+y_0^2}{\Omega(t)}} \left(\cosh\left(\frac{2yy_0}{\Omega(t)}\right) - \sinh\left(\frac{2yy_0}{\Omega(t)}\right) \right), \end{aligned} \quad (13)$$

where $\Omega(t) = \epsilon^2 + 2i\hbar t/m$.

In the usual case of a two-slit experiment, the amplitudes coming from the two slits are approximately equal, i.e., $a = b = 1/\sqrt{2}$. In this case, the sinh terms cancel out to give the dark fringes, and what is left is

$$\begin{aligned} \psi(y, t) &= \frac{1}{2}aC(t) \left(e^{-\frac{(y-y_0)^2}{\Omega(t)}} + e^{-\frac{(y+y_0)^2}{\Omega(t)}} \right) \\ &+ \frac{1}{2}bC(t) \left(e^{-\frac{(y-y_0)^2}{\Omega(t)}} + e^{-\frac{(y+y_0)^2}{\Omega(t)}} \right). \end{aligned} \quad (14)$$

In this state, the parts of the state coming from the two slits are equal. So, there is no which-way information in the state any more. If a lens is used after this stage, which takes the part $e^{-(y-y_0)^2/\Omega(t)}$ to one detector and the part $e^{-(y+y_0)^2/\Omega(t)}$ to the other detector, one can easily see from (14) that a part coming from one slit, becomes a superposition of two parts going to the two detectors. So, each detectors receives equal contribution from the two slits, and registering a particle gives no information on which slit the particle came from.

Note however, that if one of the slits is closed, say, if $a = 1$ and $b = 0$, the state at time t will be $aC(t)e^{-\frac{(y-y_0)^2}{\Omega(t)}}$, which goes to only one detector when a lens is used.




FIG. 2: A schematic representation of Wheeler's experiment¹². A photon beam is split at a half-silvered mirror S . One part of the beam goes and reflects off the mirror A and other reflects off mirror B . The beams cross and there after the detectors D_A and D_B collect the photons.

A critic might argue that instead of canceling out some parts of the two wave-packets, one might just evolve them separately, and because they were initially orthogonal, they will give distinct result at the end. The answer to that is, if one really looks for interference by putting thin wires, one is blocking those very parts of the wave-packets which are canceling out. So, those parts will not reach the detectors, even if one insists on not canceling them out. On the other hand, the argument of bringing in wires is not really needed. The state (14) as such, has no which path information.

At this stage we would like to recall an experiment discussed by Wheeler to demonstrate complementarity¹². The scheme of the experiment is shown in FIG. 2. A photon beam is split at a half-silvered mirror S . One part of the beam goes and reflects off the mirror A and other reflects off mirror B . The beams cross and there after the detectors D_A and D_B collect the photons. Interestingly, Wheeler assumes that when the two beams

cross uninterrupted, the detectors D_A and D_B give the which path information. From our preceding analysis it is clear that if there is interference in the crossing region, the two detectors no longer give the which-path information. Wheeler was contrasting this setup with the one where in the crossing region there is another half silvered mirror which mixes the two beams. Wheeler probably did not look into the possibility of an interference in the beam-crossing region. In a general case of the crossing of beams, there will be interference, and consequently, the which-path information will be lost.

In conclusion, although Afshar's experiment does have genuine interference, as shown by his introducing thin wires, the detectors detecting the photons behind the converging lens, do not yield any which-way information. Many earlier analysis of complementarity have concentrated on showing how existence of which-way information destroys interference. We have taken the reverse approach, as demanded by the Afshar experiment. We have shown that the existence of interference necessarily destroys the which-way information. The complementarity principle is robust and cannot be violated in any experiment which is a variant of the two-slit experiment. We see yet again that in dealing with quantum systems, trusting classical intuition can easily lead one to wrong conclusions.

Acknowledgments

The author thanks the Centre for Philosophy and Foundations of Science, New Delhi, for organizing annual meetings on foundations of quantum mechanics, which provide a platform for exchange of ideas.

* Electronic address: tabish@jamia-physics.net

¹ N. Bohr in *Albert Einstein philosopher-scientist*, edited by P. A. Schilpp, (The library of living philosophers, Evanston, 1949), pp. 200-241.

² M. O. Scully and K. Drühl, "Quantum eraser: A proposed photon correlation experiment concerning observation and delayed choice in quantum mechanics", *Phys. Rev. A* **25**, 2208 (1982).

³ M. O. Scully, B.-G. Englert and H. Walther, "Quantum optical tests of complementarity", *Nature (London)* **351**, 111 (1991).

⁴ "Quantum rebel" M. Chown, *New Scientist* **183**, 30 (2004).

⁵ "Violation of the principle of Complementarity, and its implications" S. S. Afshar, *Proc. SPIE* **5866** (2005) 229-244.

⁶ P.G. Kwiat, H. Weinfurter, T. Herzog, A. Zeilinger, and M.A. Kasevich, "Interaction-free measurement," *Phys. Rev. Lett.* **74**, (1995) 4763.

⁷ "Why the Afshar Experiment Does Not Refute Complementarity" R. E. Kastner, arXiv.org:quant-ph/0502021.

⁸ "Logical analysis of the Bohr Complementarity Principle in Afshar's experiment under the NAFL interpretation" R. Srinivasan, arXiv.org:quant-ph/0504115.

⁹ "Complementarity and Afshar's experiment" A. Drezet, arXiv.org:quant-ph/0508091.

¹⁰ "Afshar's Experiment does not show a Violation of Complementarity" O. Steuernagel, arXiv.org:quant-ph/0512123.

¹¹ "Entanglement and quantum interference" P. O'Hara, arXiv.org:quant-ph/0608202.

¹² "Law without law J. A. Wheeler, in *Quantum Theory and Measurement* (ed. J.A. Wheeler and W.H. Zurek), Princeton University Press pp. 182-213 (1983).